

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200407)130:1;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- Rapid communications
- Commentary
- Special issue proposals are also considered
- New initiatives—a new format for rapid publication
- Short contributions
- Letters to the Editor
- Reviews

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200407)130:1;1-1

02022

Volume 130 No. 2 July 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 130 Number 2 2004

- 135 Morphology, chemistry and distribution of neofomed spherulites in agricultural land affected by metallurgical point-source pollution
S. Leguédais, F. van Oort, T. Jongmans, P. Chevallier
Metal spherulites may act as high metal-trapping mineral phases in polluted agricultural soils.
- 149 Assessment of macro and microelement accumulation capability of two aquatic plants
D. Baldantoni, A. Alfani, P. Di Tommasi, G. Bartoli, A. Virzo De Santo
Element accumulation in roots and shoots of aquatic plants was used as a criteria for selecting useful biomonitors.
- 157 Effect of anions and cations on cadmium sorption kinetics from aqueous solutions by chitin: experimental studies and modeling
H. Benaissa, B. Benguella
Various ions can influence the ability of chitin to remove cadmium from aqueous solutions.
- 165 Trichloroacetic acid cycling in Sitka spruce saplings and effects on sapling health following long term exposure
C.A. Dickey, K.V. Heal, R.T. Stidson, R. Koren, P. Schröder, J.N. Cape, M.R. Heal
TCA stored in Sitka spruce needles may affect the health of saplings.
- 177 Organotin compounds in precipitation, fog and soils of a forested ecosystem in Germany
J.-H. Huang, D. Schwesig, E. Matzner
Forested soils may act as sinks for atmospherically deposited organotin compounds.
- 187 Polybrominated diphenyl ethers (PBDEs) and their methoxylated derivatives in pike from Swedish waters with emphasis on temporal trends, 1967–2000
A. Kierkegaard, A. Bignert, U. Sellström, M. Olsson, L. Asplund, B. Jansson, C.A. de Wit
PBDEs and MeO-BDEs show different temporal trends in pike from a Swedish lake suggesting different sources.
- 199 Responses of hybrid poplar clones and red maple seedlings to ambient O₃ under differing light within a mixed hardwood forest
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Under natural forest conditions and ambient O₃ exposures, available light plays a significant role in determining O₃ uptake and resulting physiological changes leading to the development of foliar symptoms.
- 215 Influence of light fleck and low light on foliar injury and physiological responses of two hybrid poplar clones to ozone
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Intermittent light flecks enhance physiological injury leading to increased foliar symptoms in clones of ozone-sensitive hybrid poplar.
- 229 Tracing sewage pollution using linear alkylbenzenes (LABs) in surface sediments at the south end of the Southern California Bight
J.V. Macías-Zamora, N. Ramirez-Alvarez
Linear alkylbenzenes were used to trace the distribution of sewage pollution of surface marine sediments.
- 239 Sediment trace metal profiles in lakes of Killarney Park, Canada from regional to continental influence
N. Belzile, Y.-W. Chen, J.M. Gunn, S.S. Dixit
Lakes in Killarney Park are in transition from being impacted by regional pollution to being controlled by continental atmospheric deposition.
- 249 Occurrence of PAHs, PCBs and organochlorine pesticides in the Tonghui River of Beijing, China
Z. Zhang, J. Huang, G. Yu, H. Hong
Organic pollutants were higher in sediments than in surface waters.
- 263 Macroinvertebrate response to acid mine drainage: community metrics and on-line behavioural toxicity bioassay
A. Gerhardt, L. Janssens de Bisthoven, A.M.V.M. Soares
*Toxicity of acid mine drainage was evaluated by macroinvertebrate bioassessment and a new on-line rapid behavioural toxicity test with *Atyaephyra desmaresti* (Crustacea).*

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



0269-7491(200407)130:2;1-0



ELSEVIER

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200407)130:1;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
*This is the first report of *LCT1* in Ca acquisition and regulation of Cd-toxicity by Ca.*
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200407)130:1;1-1

02022

Volume 130 No. 2 July 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 130 Number 2 2004

- 135 Morphology, chemistry and distribution of neofomed spherulites in agricultural land affected by metallurgical point-source pollution
S. Leguédais, F. van Oort, T. Jongmans, P. Chevallier
Metal spherulites may act as high metal-trapping mineral phases in polluted agricultural soils.
- 149 Assessment of macro and microelement accumulation capability of two aquatic plants
D. Baldantoni, A. Alfani, P. Di Tommasi, G. Bartoli, A. Virzo De Santo
Element accumulation in roots and shoots of aquatic plants was used as a criteria for selecting useful biomonitors.
- 157 Effect of anions and cations on cadmium sorption kinetics from aqueous solutions by chitin: experimental studies and modeling
H. Benaissa, B. Benguella
Various ions can influence the ability of chitin to remove cadmium from aqueous solutions.
- 165 Trichloroacetic acid cycling in Sitka spruce saplings and effects on sapling health following long term exposure
C.A. Dickey, K.V. Heal, R.T. Stidson, R. Koren, P. Schröder, J.N. Cape, M.R. Heal
TCA stored in Sitka spruce needles may affect the health of saplings.
- 177 Organotin compounds in precipitation, fog and soils of a forested ecosystem in Germany
J.-H. Huang, D. Schwesig, E. Matzner
Forested soils may act as sinks for atmospherically deposited organotin compounds.
- 187 Polybrominated diphenyl ethers (PBDEs) and their methoxylated derivatives in pike from Swedish waters with emphasis on temporal trends, 1967–2000
A. Kierkegaard, A. Bignert, U. Sellström, M. Olsson, L. Asplund, B. Jansson, C.A. de Wit
PBDEs and MeO-BDEs show different temporal trends in pike from a Swedish lake suggesting different sources.
- 199 Responses of hybrid poplar clones and red maple seedlings to ambient O₃ under differing light within a mixed hardwood forest
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Under natural forest conditions and ambient O₃ exposures, available light plays a significant role in determining O₃ uptake and resulting physiological changes leading to the development of foliar symptoms.
- 215 Influence of light fleck and low light on foliar injury and physiological responses of two hybrid poplar clones to ozone
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Intermittent light flecks enhance physiological injury leading to increased foliar symptoms in clones of ozone-sensitive hybrid poplar.
- 229 Tracing sewage pollution using linear alkylbenzenes (LABs) in surface sediments at the south end of the Southern California Bight
J.V. Macías-Zamora, N. Ramirez-Alvarez
Linear alkylbenzenes were used to trace the distribution of sewage pollution of surface marine sediments.
- 239 Sediment trace metal profiles in lakes of Killarney Park, Canada from regional to continental influence
N. Belzile, Y.-W. Chen, J.M. Gunn, S.S. Dixit
Lakes in Killarney Park are in transition from being impacted by regional pollution to being controlled by continental atmospheric deposition.
- 249 Occurrence of PAHs, PCBs and organochlorine pesticides in the Tonghui River of Beijing, China
Z. Zhang, J. Huang, G. Yu, H. Hong
Organic pollutants were higher in sediments than in surface waters.
- 263 Macroinvertebrate response to acid mine drainage: community metrics and on-line behavioural toxicity bioassay
A. Gerhardt, L. Janssens de Bisthoven, A.M.V.M. Soares
*Toxicity of acid mine drainage was evaluated by macroinvertebrate bioassessment and a new on-line rapid behavioural toxicity test with *Atyaephyra desmaresti* (Crustacea).*

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




0269-7491(200407)130:2;1-0



ELSEVIER

02022

Volume 130 No. 3 August 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 130 Number 3 2004

- 301 Effects of incubation on solubility and mobility of trace metals in two contaminated soils
L.Q. Ma, Y. Dong
Iron is important in controlling metal solubility and mobility in flooded soils.
- 309 Multi-generation cadmium acclimation and tolerance in *Daphnia magna* Straus
B.T.A. Muysen, C.R. Janssen
Multi-generation acclimation studies are important for evaluating long-term effects of aquatic pollutants.
- 317 The role of glomalin, a protein produced by arbuscular mycorrhizal fungi, in sequestering potentially toxic elements
M.C. González-Chávez, R. Carrillo-González, S.F. Wright, K.A. Nichols
Glomalin may be useful in remediation of toxic elements in soils.
- 325 Natural attenuation of trichloroethene and its degradation products at a lake-shore site
Y.-J. An, D.H. Kampbell, J.W. Weaver, J.T. Wilson, S.-W. Jeong
Field investigation of TCE contamination at a lake-shore site indicates that TCE is anaerobically degrading under ambient conditions.
- 337 Long-term effects of submergence and wetland vegetation on metals in a 90-year old abandoned Pb-Zn mine tailings pond
D.L. Jacob, M.L. Otte
Metal and acid-volatile sulfide concentrations were affected differently by flooding vegetation.
- 347 Contamination of butyltin compounds in Malaysian marine environments
A. Sudaryanto, S. Takahashi, H. Iwata, S. Tanabe, A. Ismail
Lack of any regulation of TBT clearly resulted in a heavy contamination of BTs in Malaysia.
- 359 The influence of season and leaf age on concentrations of radiocaesium (¹³⁷Cs), stable caesium (¹³³Cs) and potassium in *Agrostis capillaris*
C.A. Salt, J.W. Kay, K.E. Jarvis
*Season and leaf age have the same influence on radiocaesium and stable caesium in *Agrostis capillaris* but affect K differently.*
- 371 Haloacetic acids in the aquatic environment. Part I: macrophyte toxicity
M.L. Hanson, K.R. Solomon
Monochloroacetic acid was the most toxic form to aquatic macrophytes.
- 385 Haloacetic acids in the aquatic environment. Part II: ecological risk assessment
M.L. Hanson, K.R. Solomon
Current environmental concentrations of haloacetic acids do not pose a risk to aquatic macrophytes, but could impact plankton.
- 403 Comparing aquatic risk assessment methods for the photosynthesis-inhibiting herbicides metribuzin and metamiton
T.C.M. Brock, S.J.H. Crum, J.W. Deneer, F. Heimbach, R.M.M. Roijackers, J.A. Sinkeldam
Laboratory toxicity data for aquatic primary producers can be used to derive ecological thresholds for herbicides.
- 427 Arbuscular mycorrhizal fungi can decrease the uptake of uranium by subterranean clover grown at high levels of uranium in soil
G. Ruyfikiri, L. Huysmans, J. Wannijn, M. Van Hees, C. Leyval, I. Jakobsen
Plant mycorrhization may decrease U concentration in shoots of plants grown at high level of U in soil.
- 437 Degradation of toxaphene in water during anaerobic and aerobic conditions
M. LacayoR, B. van Bavel, B. Mattiasson
The majority of toxaphene degradation occurred under anaerobic conditions.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; ISI GeoSciTech; Science Citation Index; SciSearch)



0269-7491(200408)130:3;1-Z



ELSEVIER

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- Rapid communications
- Commentary
- Special issue proposals are also considered
- New initiatives—a new format for rapid publication
- Short contributions
- Letters to the Editor
- Reviews

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGÜREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
*This is the first report of *LCT1* in Ca acquisition and regulation of Cd-toxicity by Ca.*
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200407)130:1;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200407)130:1;1-1

02022

Volume 130 No. 2 July 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 130 Number 2 2004

- 135 Morphology, chemistry and distribution of neofomed spherulites in agricultural land affected by metallurgical point-source pollution
S. Leguédais, F. van Oort, T. Jongmans, P. Chevallier
Metal spherulites may act as high metal-trapping mineral phases in polluted agricultural soils.
- 149 Assessment of macro and microelement accumulation capability of two aquatic plants
D. Baldantoni, A. Alfani, P. Di Tommasi, G. Bartoli, A. Virzo De Santo
Element accumulation in roots and shoots of aquatic plants was used as a criteria for selecting useful biomonitors.
- 157 Effect of anions and cations on cadmium sorption kinetics from aqueous solutions by chitin: experimental studies and modeling
H. Benaissa, B. Benguella
Various ions can influence the ability of chitin to remove cadmium from aqueous solutions.
- 165 Trichloroacetic acid cycling in Sitka spruce saplings and effects on sapling health following long term exposure
C.A. Dickey, K.V. Heal, R.T. Stidson, R. Koren, P. Schröder, J.N. Cape, M.R. Heal
TCA stored in Sitka spruce needles may affect the health of saplings.
- 177 Organotin compounds in precipitation, fog and soils of a forested ecosystem in Germany
J.-H. Huang, D. Schwesig, E. Matzner
Forested soils may act as sinks for atmospherically deposited organotin compounds.
- 187 Polybrominated diphenyl ethers (PBDEs) and their methoxylated derivatives in pike from Swedish waters with emphasis on temporal trends, 1967–2000
A. Kierkegaard, A. Bignert, U. Sellström, M. Olsson, L. Asplund, B. Jansson, C.A. de Wit
PBDEs and MeO-BDEs show different temporal trends in pike from a Swedish lake suggesting different sources.
- 199 Responses of hybrid poplar clones and red maple seedlings to ambient O₃ under differing light within a mixed hardwood forest
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Under natural forest conditions and ambient O₃ exposures, available light plays a significant role in determining O₃ uptake and resulting physiological changes leading to the development of foliar symptoms.
- 215 Influence of light fleck and low light on foliar injury and physiological responses of two hybrid poplar clones to ozone
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Intermittent light flecks enhance physiological injury leading to increased foliar symptoms in clones of ozone-sensitive hybrid poplar.
- 229 Tracing sewage pollution using linear alkylbenzenes (LABs) in surface sediments at the south end of the Southern California Bight
J.V. Macías-Zamora, N. Ramirez-Alvarez
Linear alkylbenzenes were used to trace the distribution of sewage pollution of surface marine sediments.
- 239 Sediment trace metal profiles in lakes of Killarney Park, Canada from regional to continental influence
N. Belzile, Y.-W. Chen, J.M. Gunn, S.S. Dixit
Lakes in Killarney Park are in transition from being impacted by regional pollution to being controlled by continental atmospheric deposition.
- 249 Occurrence of PAHs, PCBs and organochlorine pesticides in the Tonghui River of Beijing, China
Z. Zhang, J. Huang, G. Yu, H. Hong
Organic pollutants were higher in sediments than in surface waters.
- 263 Macroinvertebrate response to acid mine drainage: community metrics and on-line behavioural toxicity bioassay
A. Gerhardt, L. Janssens de Bisthoven, A.M.V.M. Soares
*Toxicity of acid mine drainage was evaluated by macroinvertebrate bioassessment and a new on-line rapid behavioural toxicity test with *Atyaephyra desmaresti* (Crustacea).*

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



0269-7491(200407)130:2;1-0



ELSEVIER

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
*This is the first report of *LCT1* in Ca acquisition and regulation of Cd-toxicity by Ca.*
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
*This is the first report of *LCT1* in Ca acquisition and regulation of Cd-toxicity by Ca.*
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200407)130:1;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
*This is the first report of *LCT1* in Ca acquisition and regulation of Cd-toxicity by Ca.*
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200407)130:1;1-1

02022

Volume 130 No. 2 July 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 130 Number 2 2004

- 135 Morphology, chemistry and distribution of neofomed spherulites in agricultural land affected by metallurgical point-source pollution
S. Leguédais, F. van Oort, T. Jongmans, P. Chevallier
Metal spherulites may act as high metal-trapping mineral phases in polluted agricultural soils.
- 149 Assessment of macro and microelement accumulation capability of two aquatic plants
D. Baldantoni, A. Alfani, P. Di Tommasi, G. Bartoli, A. Virzo De Santo
Element accumulation in roots and shoots of aquatic plants was used as a criteria for selecting useful biomonitors.
- 157 Effect of anions and cations on cadmium sorption kinetics from aqueous solutions by chitin: experimental studies and modeling
H. Benaissa, B. Benguella
Various ions can influence the ability of chitin to remove cadmium from aqueous solutions.
- 165 Trichloroacetic acid cycling in Sitka spruce saplings and effects on sapling health following long term exposure
C.A. Dickey, K.V. Heal, R.T. Stidson, R. Koren, P. Schröder, J.N. Cape, M.R. Heal
TCA stored in Sitka spruce needles may affect the health of saplings.
- 177 Organotin compounds in precipitation, fog and soils of a forested ecosystem in Germany
J.-H. Huang, D. Schwesig, E. Matzner
Forested soils may act as sinks for atmospherically deposited organotin compounds.
- 187 Polybrominated diphenyl ethers (PBDEs) and their methoxylated derivatives in pike from Swedish waters with emphasis on temporal trends, 1967–2000
A. Kierkegaard, A. Bignert, U. Sellström, M. Olsson, L. Asplund, B. Jansson, C.A. de Wit
PBDEs and MeO-BDEs show different temporal trends in pike from a Swedish lake suggesting different sources.
- 199 Responses of hybrid poplar clones and red maple seedlings to ambient O₃ under differing light within a mixed hardwood forest
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Under natural forest conditions and ambient O₃ exposures, available light plays a significant role in determining O₃ uptake and resulting physiological changes leading to the development of foliar symptoms.
- 215 Influence of light fleck and low light on foliar injury and physiological responses of two hybrid poplar clones to ozone
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Intermittent light flecks enhance physiological injury leading to increased foliar symptoms in clones of ozone-sensitive hybrid poplar.
- 229 Tracing sewage pollution using linear alkylbenzenes (LABs) in surface sediments at the south end of the Southern California Bight
J.V. Macías-Zamora, N. Ramirez-Alvarez
Linear alkylbenzenes were used to trace the distribution of sewage pollution of surface marine sediments.
- 239 Sediment trace metal profiles in lakes of Killarney Park, Canada from regional to continental influence
N. Belzile, Y.-W. Chen, J.M. Gunn, S.S. Dixit
Lakes in Killarney Park are in transition from being impacted by regional pollution to being controlled by continental atmospheric deposition.
- 249 Occurrence of PAHs, PCBs and organochlorine pesticides in the Tonghui River of Beijing, China
Z. Zhang, J. Huang, G. Yu, H. Hong
Organic pollutants were higher in sediments than in surface waters.
- 263 Macroinvertebrate response to acid mine drainage: community metrics and on-line behavioural toxicity bioassay
A. Gerhardt, L. Janssens de Bisthoven, A.M.V.M. Soares
*Toxicity of acid mine drainage was evaluated by macroinvertebrate bioassessment and a new on-line rapid behavioural toxicity test with *Atyaephyra desmaresti* (Crustacea).*

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




0269-7491(200407)130:2;1-0



ELSEVIER

02022

Volume 130 No. 3 August 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 130 Number 3 2004

- 301 Effects of incubation on solubility and mobility of trace metals in two contaminated soils
L.Q. Ma, Y. Dong
Iron is important in controlling metal solubility and mobility in flooded soils.
- 309 Multi-generation cadmium acclimation and tolerance in *Daphnia magna* Straus
B.T.A. Muysen, C.R. Janssen
Multi-generation acclimation studies are important for evaluating long-term effects of aquatic pollutants.
- 317 The role of glomalin, a protein produced by arbuscular mycorrhizal fungi, in sequestering potentially toxic elements
M.C. González-Chávez, R. Carrillo-González, S.F. Wright, K.A. Nichols
Glomalin may be useful in remediation of toxic elements in soils.
- 325 Natural attenuation of trichloroethene and its degradation products at a lake-shore site
Y.-J. An, D.H. Kampbell, J.W. Weaver, J.T. Wilson, S.-W. Jeong
Field investigation of TCE contamination at a lake-shore site indicates that TCE is anaerobically degrading under ambient conditions.
- 337 Long-term effects of submergence and wetland vegetation on metals in a 90-year old abandoned Pb-Zn mine tailings pond
D.L. Jacob, M.L. Otte
Metal and acid-volatile sulfide concentrations were affected differently by flooding vegetation.
- 347 Contamination of butyltin compounds in Malaysian marine environments
A. Sudaryanto, S. Takahashi, H. Iwata, S. Tanabe, A. Ismail
Lack of any regulation of TBT clearly resulted in a heavy contamination of BTs in Malaysia.
- 359 The influence of season and leaf age on concentrations of radiocaesium (¹³⁷Cs), stable caesium (¹³³Cs) and potassium in *Agrostis capillaris*
C.A. Salt, J.W. Kay, K.E. Jarvis
*Season and leaf age have the same influence on radiocaesium and stable caesium in *Agrostis capillaris* but affect K differently.*
- 371 Haloacetic acids in the aquatic environment. Part I: macrophyte toxicity
M.L. Hanson, K.R. Solomon
Monochloroacetic acid was the most toxic form to aquatic macrophytes.
- 385 Haloacetic acids in the aquatic environment. Part II: ecological risk assessment
M.L. Hanson, K.R. Solomon
Current environmental concentrations of haloacetic acids do not pose a risk to aquatic macrophytes, but could impact plankton.
- 403 Comparing aquatic risk assessment methods for the photosynthesis-inhibiting herbicides metribuzin and metamitron
T.C.M. Brock, S.J.H. Crum, J.W. Deneer, F. Heimbach, R.M.M. Roijackers, J.A. Sinkeldam
Laboratory toxicity data for aquatic primary producers can be used to derive ecological thresholds for herbicides.
- 427 Arbuscular mycorrhizal fungi can decrease the uptake of uranium by subterranean clover grown at high levels of uranium in soil
G. Ruyfikiri, L. Huysmans, J. Wannijn, M. Van Hees, C. Leyval, I. Jakobsen
Plant mycorrhization may decrease U concentration in shoots of plants grown at high level of U in soil.
- 437 Degradation of toxaphene in water during anaerobic and aerobic conditions
M. LacayoR, B. van Bavel, B. Mattiasson
The majority of toxaphene degradation occurred under anaerobic conditions.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; ISI GeoSciTech; Science Citation Index; SciSearch)




0269-7491(200408)130:3;1-Z



ELSEVIER

02022

Volume 131 No. 1 September 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 131 Number 1 2004

- 1 Intra-specific variability of ozone sensitivity in *Centaurea jacea* L., a potential bioindicator for elevated ozone concentrations**
S. Bassin, R. Kölliker, C. Cretton, M. Bertossa, F. Widmer, P. Bungener, J. Fuhrer
*Ozone sensitivity in genetically different brown knapweed (*Centaurea jacea* L.) populations from five European countries is related to phenological development.*
- 13 Polycyclic aromatic hydrocarbon composition in soils and sediments of high altitude lakes**
J.O. Grimalt, B.L. van Drooge, A. Ribes, P. Fernández, P. Appleby
The concentrations of polycyclic hydrocarbons in soils are more strongly correlated with deposition rate than the concentrations in sediments.
- 25 Effects of invertebrate predators and a pesticide on temporary pond microcosms used for aquatic toxicity testing**
M.J. Barry, W. Davies
This study demonstrates that the use of temporary pond microcosms containing invertebrate predators may be a robust and useful method for ecosystem level toxicity testing.
- 35 Concentrations and gas/particle partitioning of PCBs in Chicago**
Y. Tasdemir, N. Vardar, M. Odabasi, T.M. Holsen
Results suggest that the area southwest of Chicago can be an important local or regional source sector for PCBs.
- 45 Treatment of atrazine by integrating photocatalytic and biological processes**
C.Y. Chan, S. Tao, R. Dawson, P.K. Wong
Photochemical pretreatment plus microbial degradation lead to complete degradation and detoxification of atrazine.
- 55 Variability of catechin and 4-hydroxyacetophenone distribution in Norway spruce needles in relation to their position, age, and growing conditions**
N. Vrhotová, J. Triska, O. Urban, L. Pěkníc
Catechin concentration in spruce needles was influenced by tree position and needle position in the tree canopy.
- 61 Solute transport and extraction by a single root in unsaturated soils: model development and experiment**
J. Kim, K. Sung, M.Y. Corapcioglu, M.C. Drew
The single-root model and experiment were useful in understanding microscopic phenomena of contaminant transport in soil-plant system.
- 71 Further investigation of the heavy metal content of the teeth of the bank vole as an exposure indicator of environmental pollution in Poland**
J. Gdula-Argasińska, J. Appleton, K. Sawicka-Kapusta, B. Spence
The molar teeth of bank voles are sensitive and accurate bio-indicators of heavy metals' exposure and reflect decreased metals deposition related to the improved environmental conditions.
- 81 A lysimeter experiment to investigate temporal changes in the availability of pesticide residues for leaching**
F.G. Renaud, C.D. Brown, C.J. Fryer, A. Walker
Leaching of aged herbicides varies with properties of the compound and of the soil.
- 93 Comparison of throughfall and soil solution chemistry between a high-density Corsican pine stand and a naturally regenerated silver birch stand**
A. De Schrijver, L. Nachtergale, J. Staelens, S. Luysaert, L. De Keersmaeker
Nitrogen throughfall deposition in a naturally regenerated birch stand was 3.5 times lower than in a neighbouring Corsican pine plantation.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; ISI GeoSciTech; Science Citation Index; SciSearch)



0269-7491(200409)131:1;1-Z



ELSEVIER

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200407)130:1;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- Rapid communications
- Commentary
- Special issue proposals are also considered
- New initiatives—a new format for rapid publication
- Short contributions
- Letters to the Editor
- Reviews

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200407)130:1;1-1

02022

Volume 130 No. 2 July 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 130 Number 2 2004

- 135 Morphology, chemistry and distribution of neofomed spherulites in agricultural land affected by metallurgical point-source pollution
S. Leguédais, F. van Oort, T. Jongmans, P. Chevallier
Metal spherulites may act as high metal-trapping mineral phases in polluted agricultural soils.
- 149 Assessment of macro and microelement accumulation capability of two aquatic plants
D. Baldantoni, A. Alfani, P. Di Tommasi, G. Bartoli, A. Virzo De Santo
Element accumulation in roots and shoots of aquatic plants was used as a criteria for selecting useful biomonitors.
- 157 Effect of anions and cations on cadmium sorption kinetics from aqueous solutions by chitin: experimental studies and modeling
H. Benaissa, B. Benguella
Various ions can influence the ability of chitin to remove cadmium from aqueous solutions.
- 165 Trichloroacetic acid cycling in Sitka spruce saplings and effects on sapling health following long term exposure
C.A. Dickey, K.V. Heal, R.T. Stidson, R. Koren, P. Schröder, J.N. Cape, M.R. Heal
TCA stored in Sitka spruce needles may affect the health of saplings.
- 177 Organotin compounds in precipitation, fog and soils of a forested ecosystem in Germany
J.-H. Huang, D. Schwesig, E. Matzner
Forested soils may act as sinks for atmospherically deposited organotin compounds.
- 187 Polybrominated diphenyl ethers (PBDEs) and their methoxylated derivatives in pike from Swedish waters with emphasis on temporal trends, 1967–2000
A. Kierkegaard, A. Bignert, U. Sellström, M. Olsson, L. Asplund, B. Jansson, C.A. de Wit
PBDEs and MeO-BDEs show different temporal trends in pike from a Swedish lake suggesting different sources.
- 199 Responses of hybrid poplar clones and red maple seedlings to ambient O₃ under differing light within a mixed hardwood forest
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Under natural forest conditions and ambient O₃ exposures, available light plays a significant role in determining O₃ uptake and resulting physiological changes leading to the development of foliar symptoms.
- 215 Influence of light fleck and low light on foliar injury and physiological responses of two hybrid poplar clones to ozone
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Intermittent light flecks enhance physiological injury leading to increased foliar symptoms in clones of ozone-sensitive hybrid poplar.
- 229 Tracing sewage pollution using linear alkylbenzenes (LABs) in surface sediments at the south end of the Southern California Bight
J.V. Macías-Zamora, N. Ramirez-Alvarez
Linear alkylbenzenes were used to trace the distribution of sewage pollution of surface marine sediments.
- 239 Sediment trace metal profiles in lakes of Killarney Park, Canada from regional to continental influence
N. Belzile, Y.-W. Chen, J.M. Gunn, S.S. Dixit
Lakes in Killarney Park are in transition from being impacted by regional pollution to being controlled by continental atmospheric deposition.
- 249 Occurrence of PAHs, PCBs and organochlorine pesticides in the Tonghui River of Beijing, China
Z. Zhang, J. Huang, G. Yu, H. Hong
Organic pollutants were higher in sediments than in surface waters.
- 263 Macroinvertebrate response to acid mine drainage: community metrics and on-line behavioural toxicity bioassay
A. Gerhardt, L. Janssens de Bisthoven, A.M.V.M. Soares
*Toxicity of acid mine drainage was evaluated by macroinvertebrate bioassessment and a new on-line rapid behavioural toxicity test with *Atyaephyra desmaresti* (Crustacea).*

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



0269-7491(200407)130:2;1-0



ELSEVIER

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
*This is the first report of *LCT1* in Ca acquisition and regulation of Cd-toxicity by Ca.*
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGÜREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
*This is the first report of *LCT1* in Ca acquisition and regulation of Cd-toxicity by Ca.*
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
*This is the first report of *LCT1* in Ca acquisition and regulation of Cd-toxicity by Ca.*
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200407)130:1;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of BauxsolTM and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined BauxsolTM and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
*This is the first report of *LCT1* in Ca acquisition and regulation of Cd-toxicity by Ca.*
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants


Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200407)130:1;1-1

02022

Volume 130 No. 2 July 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 130 Number 2 2004

- 135 Morphology, chemistry and distribution of neofomed spherulites in agricultural land affected by metallurgical point-source pollution
S. Leguédais, F. van Oort, T. Jongmans, P. Chevallier
Metal spherulites may act as high metal-trapping mineral phases in polluted agricultural soils.
- 149 Assessment of macro and microelement accumulation capability of two aquatic plants
D. Baldantoni, A. Alfani, P. Di Tommasi, G. Bartoli, A. Virzo De Santo
Element accumulation in roots and shoots of aquatic plants was used as a criteria for selecting useful biomonitors.
- 157 Effect of anions and cations on cadmium sorption kinetics from aqueous solutions by chitin: experimental studies and modeling
H. Benaissa, B. Benguella
Various ions can influence the ability of chitin to remove cadmium from aqueous solutions.
- 165 Trichloroacetic acid cycling in Sitka spruce saplings and effects on sapling health following long term exposure
C.A. Dickey, K.V. Heal, R.T. Stidson, R. Koren, P. Schröder, J.N. Cape, M.R. Heal
TCA stored in Sitka spruce needles may affect the health of saplings.
- 177 Organotin compounds in precipitation, fog and soils of a forested ecosystem in Germany
J.-H. Huang, D. Schwesig, E. Matzner
Forested soils may act as sinks for atmospherically deposited organotin compounds.
- 187 Polybrominated diphenyl ethers (PBDEs) and their methoxylated derivatives in pike from Swedish waters with emphasis on temporal trends, 1967–2000
A. Kierkegaard, A. Bignert, U. Sellström, M. Olsson, L. Asplund, B. Jansson, C.A. de Wit
PBDEs and MeO-BDEs show different temporal trends in pike from a Swedish lake suggesting different sources.
- 199 Responses of hybrid poplar clones and red maple seedlings to ambient O₃ under differing light within a mixed hardwood forest
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Under natural forest conditions and ambient O₃ exposures, available light plays a significant role in determining O₃ uptake and resulting physiological changes leading to the development of foliar symptoms.
- 215 Influence of light fleck and low light on foliar injury and physiological responses of two hybrid poplar clones to ozone
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Intermittent light flecks enhance physiological injury leading to increased foliar symptoms in clones of ozone-sensitive hybrid poplar.
- 229 Tracing sewage pollution using linear alkylbenzenes (LABs) in surface sediments at the south end of the Southern California Bight
J.V. Macías-Zamora, N. Ramirez-Alvarez
Linear alkylbenzenes were used to trace the distribution of sewage pollution of surface marine sediments.
- 239 Sediment trace metal profiles in lakes of Killarney Park, Canada from regional to continental influence
N. Belzile, Y.-W. Chen, J.M. Gunn, S.S. Dixit
Lakes in Killarney Park are in transition from being impacted by regional pollution to being controlled by continental atmospheric deposition.
- 249 Occurrence of PAHs, PCBs and organochlorine pesticides in the Tonghui River of Beijing, China
Z. Zhang, J. Huang, G. Yu, H. Hong
Organic pollutants were higher in sediments than in surface waters.
- 263 Macroinvertebrate response to acid mine drainage: community metrics and on-line behavioural toxicity bioassay
A. Gerhardt, L. Janssens de Bisthoven, A.M.V.M. Soares
*Toxicity of acid mine drainage was evaluated by macroinvertebrate bioassessment and a new on-line rapid behavioural toxicity test with *Atyaephyra desmaresti* (Crustacea).*

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




0269-7491(200407)130:2;1-0



ELSEVIER

02022

Volume 130 No. 3 August 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 130 Number 3 2004

- 301 Effects of incubation on solubility and mobility of trace metals in two contaminated soils
L.Q. Ma, Y. Dong
Iron is important in controlling metal solubility and mobility in flooded soils.
- 309 Multi-generation cadmium acclimation and tolerance in *Daphnia magna* Straus
B.T.A. Muysen, C.R. Janssen
Multi-generation acclimation studies are important for evaluating long-term effects of aquatic pollutants.
- 317 The role of glomalin, a protein produced by arbuscular mycorrhizal fungi, in sequestering potentially toxic elements
M.C. González-Chávez, R. Carrillo-González, S.F. Wright, K.A. Nichols
Glomalin may be useful in remediation of toxic elements in soils.
- 325 Natural attenuation of trichloroethene and its degradation products at a lake-shore site
Y.-J. An, D.H. Kampbell, J.W. Weaver, J.T. Wilson, S.-W. Jeong
Field investigation of TCE contamination at a lake-shore site indicates that TCE is anaerobically degrading under ambient conditions.
- 337 Long-term effects of submergence and wetland vegetation on metals in a 90-year old abandoned Pb-Zn mine tailings pond
D.L. Jacob, M.L. Otte
Metal and acid-volatile sulfide concentrations were affected differently by flooding vegetation.
- 347 Contamination of butyltin compounds in Malaysian marine environments
A. Sudaryanto, S. Takahashi, H. Iwata, S. Tanabe, A. Ismail
Lack of any regulation of TBT clearly resulted in a heavy contamination of BTs in Malaysia.
- 359 The influence of season and leaf age on concentrations of radiocaesium (¹³⁷Cs), stable caesium (¹³³Cs) and potassium in *Agrostis capillaris*
C.A. Salt, J.W. Kay, K.E. Jarvis
*Season and leaf age have the same influence on radiocaesium and stable caesium in *Agrostis capillaris* but affect K differently.*
- 371 Haloacetic acids in the aquatic environment. Part I: macrophyte toxicity
M.L. Hanson, K.R. Solomon
Monochloroacetic acid was the most toxic form to aquatic macrophytes.
- 385 Haloacetic acids in the aquatic environment. Part II: ecological risk assessment
M.L. Hanson, K.R. Solomon
Current environmental concentrations of haloacetic acids do not pose a risk to aquatic macrophytes, but could impact plankton.
- 403 Comparing aquatic risk assessment methods for the photosynthesis-inhibiting herbicides metribuzin and metamitron
T.C.M. Brock, S.J.H. Crum, J.W. Deneer, F. Heimbach, R.M.M. Roijackers, J.A. Sinkeldam
Laboratory toxicity data for aquatic primary producers can be used to derive ecological thresholds for herbicides.
- 427 Arbuscular mycorrhizal fungi can decrease the uptake of uranium by subterranean clover grown at high levels of uranium in soil
G. Ruyfikiri, L. Huysmans, J. Wannijn, M. Van Hees, C. Leyval, I. Jakobsen
Plant mycorrhization may decrease U concentration in shoots of plants grown at high level of U in soil.
- 437 Degradation of toxaphene in water during anaerobic and aerobic conditions
M. LacayoR, B. van Bavel, B. Mattiasson
The majority of toxaphene degradation occurred under anaerobic conditions.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; ISI GeoSciTech; Science Citation Index; SciSearch)



0269-7491(200408)130:3;1-Z



ELSEVIER

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200407)130:1;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGÜREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200407)130:1;1-1

02022

Volume 130 No. 2 July 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 130 Number 2 2004

- 135 Morphology, chemistry and distribution of neofomed spherulites in agricultural land affected by metallurgical point-source pollution
S. Leguédais, F. van Oort, T. Jongmans, P. Chevallier
Metal spherulites may act as high metal-trapping mineral phases in polluted agricultural soils.
- 149 Assessment of macro and microelement accumulation capability of two aquatic plants
D. Baldantoni, A. Alfani, P. Di Tommasi, G. Bartoli, A. Virzo De Santo
Element accumulation in roots and shoots of aquatic plants was used as a criteria for selecting useful biomonitors.
- 157 Effect of anions and cations on cadmium sorption kinetics from aqueous solutions by chitin: experimental studies and modeling
H. Benaissa, B. Benguella
Various ions can influence the ability of chitin to remove cadmium from aqueous solutions.
- 165 Trichloroacetic acid cycling in Sitka spruce saplings and effects on sapling health following long term exposure
C.A. Dickey, K.V. Heal, R.T. Stidson, R. Koren, P. Schröder, J.N. Cape, M.R. Heal
TCA stored in Sitka spruce needles may affect the health of saplings.
- 177 Organotin compounds in precipitation, fog and soils of a forested ecosystem in Germany
J.-H. Huang, D. Schwesig, E. Matzner
Forested soils may act as sinks for atmospherically deposited organotin compounds.
- 187 Polybrominated diphenyl ethers (PBDEs) and their methoxylated derivatives in pike from Swedish waters with emphasis on temporal trends, 1967–2000
A. Kierkegaard, A. Bignert, U. Sellström, M. Olsson, L. Asplund, B. Jansson, C.A. de Wit
PBDEs and MeO-BDEs show different temporal trends in pike from a Swedish lake suggesting different sources.
- 199 Responses of hybrid poplar clones and red maple seedlings to ambient O₃ under differing light within a mixed hardwood forest
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Under natural forest conditions and ambient O₃ exposures, available light plays a significant role in determining O₃ uptake and resulting physiological changes leading to the development of foliar symptoms.
- 215 Influence of light fleck and low light on foliar injury and physiological responses of two hybrid poplar clones to ozone
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Intermittent light flecks enhance physiological injury leading to increased foliar symptoms in clones of ozone-sensitive hybrid poplar.
- 229 Tracing sewage pollution using linear alkylbenzenes (LABs) in surface sediments at the south end of the Southern California Bight
J.V. Macías-Zamora, N. Ramirez-Alvarez
Linear alkylbenzenes were used to trace the distribution of sewage pollution of surface marine sediments.
- 239 Sediment trace metal profiles in lakes of Killarney Park, Canada from regional to continental influence
N. Belzile, Y.-W. Chen, J.M. Gunn, S.S. Dixit
Lakes in Killarney Park are in transition from being impacted by regional pollution to being controlled by continental atmospheric deposition.
- 249 Occurrence of PAHs, PCBs and organochlorine pesticides in the Tonghui River of Beijing, China
Z. Zhang, J. Huang, G. Yu, H. Hong
Organic pollutants were higher in sediments than in surface waters.
- 263 Macroinvertebrate response to acid mine drainage: community metrics and on-line behavioural toxicity bioassay
A. Gerhardt, L. Janssens de Bisthoven, A.M.V.M. Soares
*Toxicity of acid mine drainage was evaluated by macroinvertebrate bioassessment and a new on-line rapid behavioural toxicity test with *Atyaephyra desmaresti* (Crustacea).*

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



0269-7491(200407)130:2;1-0



ELSEVIER

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
*This is the first report of *LCT1* in Ca acquisition and regulation of Cd-toxicity by Ca.*
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200407)130:1;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants



Guest Editors
K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors


K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
*This is the first report of *LCT1* in Ca acquisition and regulation of Cd-toxicity by Ca.*
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGÜREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRRENBARGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200406)129:3;1-V

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at:
<http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200401)127:2;1-2

02022

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200402)127:3;1-0

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>


ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman**



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVA, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 1 2004

- 1 Global pollution monitoring of butyltin compounds using skipjack tuna as a bioindicator
D. UENO, S. INOUE, S. TAKAHASHI, K. IKEDA, H. TANAKA (Japan), A.N. SUBRAMANIAN (India), G. FILLMANN (Brazil), P.K.S. LAM, J. ZHENG (Hong Kong), M. MUCHTAR (Indonesia), M. PRUDENTE (Philippines), K. CHUNG (South Korea), S. TANABE (Japan)
Global pollution monitoring of butyltin in offshore water and open sea were conducted using skipjack tuna as a bioindicator.
- 13 Mutual effects of cadmium and phosphate on their adsorption and desorption by goethite
K. WANG, B. XING (USA)
Cd and phosphate have mutual influences on their sorption kinetics.
- 21 Soil ecotoxicity assessment using cadmium sensitive plants
Y.-J. AN (South Korea)
The crop plants, sorghum and cucumber, can be used as indicator species to assess ecotoxicity of soils contaminated by cadmium.
- 27 Foraminiferal proxies for pollution monitoring in moderately polluted harbors
E. ARMYNOT DU CHÂTELET, J.-P. DEBENAY, R. SOULARD (France)
Foraminiferal density and species richness that decrease with an increase in heavy metal and PAH concentration may be used as pollution indicators.
- 41 Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of ortho-hydroxychlorobiphenyls formed from PCBs by plants
K. FRANCOVA, M. MACKOVÁ, T. MACEK (Czech Republic), M. SYLVESTRE (Canada)
Bacterial dioxygenases are useful in breakdown of PCB products associated with plants.
- 49 Preliminary study of prairies forested with *Eucalyptus* sp. at the northwestern Uruguayan soils
L. CARRASCO-LETELIER (Chile), G. EGUREN, C. CASTIÑEIRA (Uruguay), O. PARRA (Chile), D. PANARIO (Uruguay)
The forestation of Uruguayan natural prairie soil does not always ensure an increase of soil carbon sink.
- 57 Throughfall chemistry and canopy interactions in a Sitka spruce plantation sprayed with six different simulated polluted mist treatments
M. CHIWA (Japan), A. CROSSLEY, L.J. SHEPPARD (UK), H. SAKUGAWA (Japan), J.N. CAPE (UK)
Weak organic acids may be involved in neutralizing acidity of throughfall.
- 65 White birch (*Betula papyrifera* Marshall) foliar litter decomposition in relation to trace metal atmospheric inputs at metal-contaminated and uncontaminated sites near Sudbury, Ontario and Rouyn-Noranda, Quebec, Canada
D. JOHNSON, B. HALE (Canada)
Copper smelter emissions in Sudbury, Ontario may cause reductions in birch litter decomposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:1;1-3

CONTENTS—Continued from outside back cover

- 73 Evaluation of chemical immobilization treatments for reducing heavy metal transport in a smelter-contaminated soil
N.T. BASTA, S.L. MCGOWEN (USA)
Diammonium phosphate was the most effective treatment for immobilizing heavy metals.
- 83 Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*)
T. IKEMOTO, T. KUNITO, I. WATANABE, G. YASUNAGA, N. BABA, N. MIYAZAKI (Japan), E.A. PETROV (Russia), S. TANABE (Japan)
Hair samples may be useful in monitoring trace elements in seals.
- 99 Combined effects of copper and food on the midge *Chironomus riparius* in whole-sediment bioassays
E.M. DE HAAS, M. LÉON PAUMEN, A.A. KOELMANS, M.H.S. KRAAK (The Netherlands)
Trophic state of an ecosystem may alter risk of sediment-bound toxicants to midges.
- 109 Regression models to predict water-soil heavy metals partition coefficients in risk assessment studies
C. CARLON (Italy), M. DALLA VALLE (UK), A. MARCOMINI (Italy)
The adoption of regression models for heavy metals could provide a substantial improvement in risk assessment practices.
- 117 Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure
S.-G. KIM, J.-H. JEE, J.-C. KANG (South Korea)
Cd accumulation and elimination in tissue of olive flounder.
- 125 High human exposure to lead through consumption of birds hunted with lead shot
P. JOHANSEN, G. ASMUND, F. RIGET (Denmark)
Lead shot contaminates the edible parts of birds so that tolerable human lead intake is exceeded.
- 131 Characterisation of a reference site for quantifying uncertainties related to soil sampling
S. BARBIZZI, P. DE ZORZI, M. BELLI, A. PATI, U. SANSONE, L. STELLATO, M. BARBINA, A. DELUISA, S. MENEGON, V. COLETTI (Italy)
An integrated approach to quality assurance in soil sampling remains to be accomplished.
- 137 Soil mobility of sewage sludge-derived dissolved organic matter, copper, nickel and zinc
D.J. ASHWORTH, B.J. ALLOWAY (UK)
Only nickel concentrations showed potential environmental significance.
- 145 Genotype-specific responses of fluctuating asymmetry and of preadult survival to the effects of lead and temperature stress in *Drosophila melanogaster*
M. POLAK, D.E. KROEGER, I.L. CARTWRIGHT, C. PONCE DELEON (USA)
Caution should be used in using fluctuating asymmetry as a stress indicator.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 2 2004

Research Papers

- 157 Effects of Bauxsol™ and biosolids on soil conditions of acid-generating mine spoil for plant growth
G. MADDOCKS, C. LIN, D. MCCONCHIE (Australia)
Application of combined Bauxsol™ and biosolids had better effects on improving conditions of acidified sulfidic mine soils for plant growth, than either alone.
- 169 Possible evidence for transport of an iron cyanide complex by plants
M. SAMIOTAKIS, S.D. EBBS (USA)
Results suggest ferrocyanide transport by barley, oat and wild cane.
- 175 Study of metal fractionation in river sediments. A comparison between kinetic and sequential extraction procedures
M.J. GISMERA, J. LACAL, P. DA SILVA, R. GARCÍA, M.T. SEVILLA, J.R. PROCOPIO (Spain)
A leached metal kinetic study may be a good way to study the mobility of trace metals in sediments.
- 183 Effects of heavy metals on population growth and metallothionein gene expression in the mosquito *Culex quinquefasciatus*, from Calcutta, India
S. SARKAR, A.K. DUTTAGUPTA (India), T.K. MAL (USA)
The mosquito, Culex quinquefasciatus, might be useful as an indicator of trace metals.
- 195 Accumulation and distribution of platinum and rhodium in the European eel *Anguilla anguilla* following aqueous exposure to metal salts
S. ZIMMERMANN, U. BAUMANN, H. TARASCHEWSKI, B. SURES (Germany)
Eels are suitable accumulation indicators for platinum group metals in aquatic ecosystems.
- 203 Mussel-based monitoring of trace metal and organic contaminants along the east coast of China using *Perna viridis* and *Mytilus edulis*
C.N. FUNG, J.C.W. LAM, G.J. ZHENG (Hong Kong), D.W. CONNELL (Australia), I. MONIRITH, S. TANABE (Japan), B.J. RICHARDSON, P.K.S. LAM (Hong Kong)
Mussel-based monitoring revealed trace metal and organic contaminants which pose potential human health risks.
- 217 Probabilistic risk assessment of abalone *Haliotis diversicolor supertexta* exposed to waterborne zinc
C.-M. LIAO, M.-P. LING (Taiwan)
A novel risk assessment method was developed for abalone.
- 229 Using sequential indicator simulation to assess the uncertainty of delineating heavy-metal contaminated soils
K.-W. JUANG, Y.-S. CHEN, D.-Y. LEE (Taiwan)
A method of geostatistical simulation was developed to demonstrate the uncertainty of assessing delineation of soils contaminated with heavy metals.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER

02022



0269-7491(200401)127:2;1-2

CONTENTS—Continued from outside back cover

- 239 Geostatistical analyses and hazard assessment on soil lead in Silvermines area, Ireland
D. MCGRATH, C. ZHANG, O.T. CARTON (Ireland)
A probability map was produced that was useful for hazard assessment and decision support.
- 249 Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies
M. FERRETTI, E. BRAMBILLA, G. BRUNIALTI, F. FORNASIER, C. MAZZALI, P. GIORDANI, P.L. NIMIS (Italy)
An acceptable level of precision must be decided before determining a sampling design.
- 257 The effects of atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming, USA—a critical review
D.A. BURNS (USA)
The effects of nitrogen deposition will become more evident as growth increases.
- 271 The possible influence of volcanic emissions on atmospheric aerosols in the city of Colima, Mexico
J. MIRANDA, F. ZEPEDA, I. GALINDO (Mexico)
Elemental analyses of PM₁₅ in the City of Colima, Mexico, were done to identify possible contributions from the Volcán de Colima, an active volcano.
- 281 Exploring field vegetation reflectance as an indicator of soil contamination in river floodplains
L. KOOISTRA, E.A.L. SALAS, J.G.P.W. CLEVERS, R. WEHRENS, R.S.E.W. LEUVEN, P.H. NIENHUIS, L.M.C. BUYDENS (The Netherlands)
Influence of elevated soil metal concentrations on field vegetation reflectance was assessed.
- 291 Seasonal effect on ammonia nitrogen removal by constructed wetlands treating polluted river water in southern Taiwan
S.-R. JING, Y.-F. LIN (Taiwan)
Seasonal variation of mass loading rate predominantly affected ammonia removal of constructed wetlands.
- 303 Effects of environmental concentrations of atrazine on hemocyte density and phagocytic activity in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. RUSSO, L. LAGADIC (France)
Phagocytosis and oxidative burst of hemocytes may be a functional biomarker for atrazine effects on snails.
- Letter to the Editor**
- 313 Regarding the rodent sperm analysis (RSA) article in the Journal of Environmental Pollution
B. TUCKER (USA)

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER

Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 127 Number 3 2004.

Research Papers

- 315 Distribution of sewage pollution around a maritime Antarctic research station indicated by faecal coliforms, *Clostridium perfringens* and faecal sterol markers
K.A. HUGHES, A. THOMPSON (UK)
Sewage contamination of seawater and marine sediments near Rothera Research Station (Antarctic Peninsula) was limited to the immediate vicinity of the outfall.
- 323 Studies of spatial and temporal distribution characteristics of TSP-bound trace metals in Seoul, Korea
K.-H. KIM, B.-J. CHOI, S.-T. YUN, S.-J. HWANG (South Korea)
Spatial and temporal factors regulate metal types differently.
- 335 The importance of biological factors affecting trace metal concentration as revealed from accumulation patterns in co-occurring terrestrial invertebrates
F. HENDRICKX, J.-P. MAELFAIT, N. BOGAERT, C. TOJAL, G. DU LAING, F.M.G. TACK, M.G. VERLOO (Belgium)
Site and species specific biological factors are important in determining contamination of biota.
- 343 Micro-spatial variation of soil metal pollution and plant recruitment near a copper smelter in Central Chile
R. GINOCCHIO, G. CARVALLO, I. TORO, E. BUSTAMANTE, Y. SILVA, N. SEPULVEDA (Chile)
Metal availability was different under shrub canopies than in open spaces.
- 353 Effects of the mosquito larvicide GB-1111 on bird eggs
D.J. HOFFMAN, P.H. ALBERS, M.J. MELANCON, A.K. MILES (USA)
Mosquito larvicide GB-1111 is unlikely to harm bobwhite embryos when applied according to product label guidance, but could pose a potential risk to mallard embryos under conditions of larvicide drift or spray overlap.
- 359 The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa
G. NAIDOO, D. CHIRKOOT (South Africa)
Coal dust reduced photosynthetic performance of the mangrove, Avicennia marina.
- 367 Effect of food level on the acute and chronic responses of daphnids to lindane
S.C. ANTUNES, B.B. CASTRO, F. GONÇALVES (Portugal)
Food level can influence responses of daphnids to lindane in toxicity tests.
- 377 Use of cyanobacteria to assess water quality in running waters
I. DOUTERELO, E. PERONA, P. MATEO (Spain)
Taxonomic composition of cyanobacteria is a sensitive indicator of river water quality.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200402)127:3;1-0

02022

CONTENTS—Continued from outside back cover

- 385 Feasibility of constructed wetlands for removing chlorothalonil and chlorpyrifos from aqueous mixtures
R.M. SHERRARD, J.S. BEARR, C.L. MURRAY-GULDE, J.H. RODGERS, JR., Y.T. SHAH (USA)
Constructed wetlands have potential for treatment of pesticide mixtures in stormwater runoff.
- 395 Biomineralisation of 1,2,4-trichlorobenzene in soils by an adapted microbial population
R. SCHROLL, F. BRAHUSHI, U. DÖRFLER, S. KÜHN (Germany), J. FEKETE (Hungary), J.C. MUNCH (Germany)
Enhanced mineralisation of 1,2,4-trichlorobenzene was observed in an agricultural soil by inoculating it with an adapted microbial community.
- 403 OH-initiated oxidation of DMS/DMSO: reaction products at high NO_x levels
V. LIBRANDO, G. TRINGALI, J. HJORTH, S. COLUCCIA (Italy)
High NO_x levels play a significant role in the atmospheric chemistry DMS/DMSO.
- 411 Organochlorine concentrations in diseased vs. healthy gull chicks from the northern Baltic
M. HARIO, J.-P. HIRVI (Finland), T. HOLLMÉN (USA), E. RUDBÄCK (Finland)
Elevated DDE/PCB ratio correlates with a high rate of chick diseases in the endangered nominate lesser black-backed gull.
- 425 Biodegradation of nonylphenol in river sediment
S.Y. YUAN, C.H. YU, B.V. CHANG (Taiwan)
The effects of manipulating several factors on nonylphenol and nonylphenol monoethoxylate degradation in river sediment were analysed.
- 431 Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment
L. XIE, P. XIE (PR China), K. OZAWA, T. HONMA, A. YOKOYAMA, H.-D. PARK (Japan)
Silver carp are tolerant of cyanobacterial toxins, and might be used to control toxic algal blooms in highly eutrophic lakes.
- 441 The use of light and electron microscopy to assess the impact of ozone on Norway spruce needles
M. KIVIMÄENPÄÄ (Finland), A.M. JÖNSSON, I. STJERNQUIST, G. SELLDÉN (Sweden), S. SUTINEN (Finland)
Light microscopy was used to diagnose ozone injury on Norway spruce needles.
- 455 Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy)
P. ADAMO, M. ARIENZO, M. PUGLIESE, V. ROCA, P. VIOLANTE (Italy)
*Elimination of radionuclides accumulated by the lichen *Stereocaulon vesuvianum* appears to be a slow process.*
- I Contents of Volume 127

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>

ENVIRONMENTAL POLLUTION

Special issue:
Persistent Organic
Pollutants

Guest Editors

K.C. Jones
A. Sweetman
D. MacKay

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



ELSEVIER



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 128 Numbers 1–2 2004

Special Issue:

Persistent Organic Pollutants

Guest Editors

K.C. Jones, A. Sweetman, D. MacKay

(For full contents see p. iii)

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200403)128:1-2;1-1

02022

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa
K.C. Jones J.W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 2 2004

Research Papers

- 165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
Vegetable-derived lubricants were more degradable than mineral oil lubricants.
- 175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.
- 183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.
- 195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRREBERGER, P. FITZE (Switzerland)
The effects of F can be seen in changes in soil chemistry, vegetation and clay mineralogy.
- 209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.
- 229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.
- 237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.
- 247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.
- 257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)



ELSEVIER



0269-7491(200405)129:2;1-X

02022

Volume 129 No. 3 June 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 129 Number 3 2004

Invited Paper

- 341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
Several alternatives are presented for poorly defined term "heavy metals".

Research Papers

- 345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
Hydrocarbon measurements cannot be substituted for actual toxicity determinations.
- 355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUJEGODA (Australia)
Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.
- 363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.
- 377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
The endpoints were analytically reproducible and sensitive parameters.
- 387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
PAH concentrations decreased with distance from the urban centre.
- 399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
A new index of coral marine pollution is proposed.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200406)129:3;1-V

02022



Volume 130 No. 1 July 2004 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

Special issue:

Effects of Air Pollution on
the Central and Eastern
European Mountain Forests

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones


J. W. Erisman

Guest Editors

A. Bytnerowicz

R. Musselman

R. Szaro



Aims and Scope:

Environmental Pollution is an international journal that addresses issues relevant to the nature, distribution and ecological effects of all types and forms of chemical pollutants in air, soil and water. The Editors welcome articles based on original research, findings from re-examination and interpretation of existing data and reviews of important issues. In addition, the journal also publishes articles on new methods of detection, study and remediation of environmental pollutants.

Types of Contributions:

- Research articles
- New initiatives—a new format for rapid publication
- Rapid communications
- Short contributions
- Reviews
- Commentary
- Letters to the Editor
- Special issue proposals are also considered

THERE ARE NO PAGE CHARGES. Twenty-five free reprints are provided.

Mission Statement: The Editors of *Environmental Pollution* seek to publish papers of the highest quality and significance and to effectively convey them to a world-wide audience in a timely manner

CONTENTS

Volume 130 Number 1 2004

Special Issue:

**Effects of Air Pollution on the Central and Eastern
European Mountain Forests**

Guest Editors

A. Bytnerowicz, R. Musselman, R. Szaro

(For full contents see p. iii)

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




ELSEVIER



0269-7491(200407)130:1;1-1

02022

Volume 130 No. 2 July 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 130 Number 2 2004

- 135 Morphology, chemistry and distribution of neofomed spherulites in agricultural land affected by metallurgical point-source pollution
S. Leguédais, F. van Oort, T. Jongmans, P. Chevallier
Metal spherulites may act as high metal-trapping mineral phases in polluted agricultural soils.
- 149 Assessment of macro and microelement accumulation capability of two aquatic plants
D. Baldantoni, A. Alfani, P. Di Tommasi, G. Bartoli, A. Virzo De Santo
Element accumulation in roots and shoots of aquatic plants was used as a criteria for selecting useful biomonitors.
- 157 Effect of anions and cations on cadmium sorption kinetics from aqueous solutions by chitin: experimental studies and modeling
H. Benaissa, B. Benguella
Various ions can influence the ability of chitin to remove cadmium from aqueous solutions.
- 165 Trichloroacetic acid cycling in Sitka spruce saplings and effects on sapling health following long term exposure
C.A. Dickey, K.V. Heal, R.T. Stidson, R. Koren, P. Schröder, J.N. Cape, M.R. Heal
TCA stored in Sitka spruce needles may affect the health of saplings.
- 177 Organotin compounds in precipitation, fog and soils of a forested ecosystem in Germany
J.-H. Huang, D. Schwesig, E. Matzner
Forested soils may act as sinks for atmospherically deposited organotin compounds.
- 187 Polybrominated diphenyl ethers (PBDEs) and their methoxylated derivatives in pike from Swedish waters with emphasis on temporal trends, 1967–2000
A. Kierkegaard, A. Bignert, U. Sellström, M. Olsson, L. Asplund, B. Jansson, C.A. de Wit
PBDEs and MeO-BDEs show different temporal trends in pike from a Swedish lake suggesting different sources.
- 199 Responses of hybrid poplar clones and red maple seedlings to ambient O₃ under differing light within a mixed hardwood forest
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Under natural forest conditions and ambient O₃ exposures, available light plays a significant role in determining O₃ uptake and resulting physiological changes leading to the development of foliar symptoms.
- 215 Influence of light fleck and low light on foliar injury and physiological responses of two hybrid poplar clones to ozone
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Intermittent light flecks enhance physiological injury leading to increased foliar symptoms in clones of ozone-sensitive hybrid poplar.
- 229 Tracing sewage pollution using linear alkylbenzenes (LABs) in surface sediments at the south end of the Southern California Bight
J.V. Macías-Zamora, N. Ramirez-Alvarez
Linear alkylbenzenes were used to trace the distribution of sewage pollution of surface marine sediments.
- 239 Sediment trace metal profiles in lakes of Killarney Park, Canada from regional to continental influence
N. Belzile, Y.-W. Chen, J.M. Gunn, S.S. Dixit
Lakes in Killarney Park are in transition from being impacted by regional pollution to being controlled by continental atmospheric deposition.
- 249 Occurrence of PAHs, PCBs and organochlorine pesticides in the Tonghui River of Beijing, China
Z. Zhang, J. Huang, G. Yu, H. Hong
Organic pollutants were higher in sediments than in surface waters.
- 263 Macroinvertebrate response to acid mine drainage: community metrics and on-line behavioural toxicity bioassay
A. Gerhardt, L. Janssens de Bisthoven, A.M.V.M. Soares
*Toxicity of acid mine drainage was evaluated by macroinvertebrate bioassessment and a new on-line rapid behavioural toxicity test with *Atyaephyra desmaresti* (Crustacea).*

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; ISI GeoSciTech; Science Citation Index; SciSearch)




0269-7491(200407)130:2;1-0



ELSEVIER

02022

Volume 130 No. 3 August 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 130 Number 3 2004

- 301 Effects of incubation on solubility and mobility of trace metals in two contaminated soils
L.Q. Ma, Y. Dong
Iron is important in controlling metal solubility and mobility in flooded soils.
- 309 Multi-generation cadmium acclimation and tolerance in *Daphnia magna* Straus
B.T.A. Muysen, C.R. Janssen
Multi-generation acclimation studies are important for evaluating long-term effects of aquatic pollutants.
- 317 The role of glomalin, a protein produced by arbuscular mycorrhizal fungi, in sequestering potentially toxic elements
M.C. González-Chávez, R. Carrillo-González, S.F. Wright, K.A. Nichols
Glomalin may be useful in remediation of toxic elements in soils.
- 325 Natural attenuation of trichloroethene and its degradation products at a lake-shore site
Y.-J. An, D.H. Kampbell, J.W. Weaver, J.T. Wilson, S.-W. Jeong
Field investigation of TCE contamination at a lake-shore site indicates that TCE is anaerobically degrading under ambient conditions.
- 337 Long-term effects of submergence and wetland vegetation on metals in a 90-year old abandoned Pb-Zn mine tailings pond
D.L. Jacob, M.L. Otte
Metal and acid-volatile sulfide concentrations were affected differently by flooding vegetation.
- 347 Contamination of butyltin compounds in Malaysian marine environments
A. Sudaryanto, S. Takahashi, H. Iwata, S. Tanabe, A. Ismail
Lack of any regulation of TBT clearly resulted in a heavy contamination of BTs in Malaysia.
- 359 The influence of season and leaf age on concentrations of radiocaesium (¹³⁷Cs), stable caesium (¹³³Cs) and potassium in *Agrostis capillaris*
C.A. Salt, J.W. Kay, K.E. Jarvis
*Season and leaf age have the same influence on radiocaesium and stable caesium in *Agrostis capillaris* but affect K differently.*
- 371 Haloacetic acids in the aquatic environment. Part I: macrophyte toxicity
M.L. Hanson, K.R. Solomon
Monochloroacetic acid was the most toxic form to aquatic macrophytes.
- 385 Haloacetic acids in the aquatic environment. Part II: ecological risk assessment
M.L. Hanson, K.R. Solomon
Current environmental concentrations of haloacetic acids do not pose a risk to aquatic macrophytes, but could impact plankton.
- 403 Comparing aquatic risk assessment methods for the photosynthesis-inhibiting herbicides metribuzin and metamitron
T.C.M. Brock, S.J.H. Crum, J.W. Deneer, F. Heimbach, R.M.M. Roijackers, J.A. Sinkeldam
Laboratory toxicity data for aquatic primary producers can be used to derive ecological thresholds for herbicides.
- 427 Arbuscular mycorrhizal fungi can decrease the uptake of uranium by subterranean clover grown at high levels of uranium in soil
G. Ruyfikiri, L. Huysmans, J. Wannijn, M. Van Hees, C. Leyval, I. Jakobsen
Plant mycorrhization may decrease U concentration in shoots of plants grown at high level of U in soil.
- 437 Degradation of toxaphene in water during anaerobic and aerobic conditions
M. LacayoR, B. van Bavel, B. Mattiasson
The majority of toxaphene degradation occurred under anaerobic conditions.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; ISI GeoSciTech; Science Citation Index; SciSearch)




0269-7491(200408)130:3;1-Z



ELSEVIER

02022

Volume 131 No. 1 September 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 131 Number 1 2004

- 1 Intra-specific variability of ozone sensitivity in *Centaurea jacea* L., a potential bioindicator for elevated ozone concentrations**
S. Bassin, R. Kölliker, C. Cretton, M. Bertossa, F. Widmer, P. Bungener, J. Fuhrer
*Ozone sensitivity in genetically different brown knapweed (*Centaurea jacea* L.) populations from five European countries is related to phenological development.*
- 13 Polycyclic aromatic hydrocarbon composition in soils and sediments of high altitude lakes**
J.O. Grimalt, B.L. van Drooge, A. Ribes, P. Fernández, P. Appleby
The concentrations of polycyclic hydrocarbons in soils are more strongly correlated with deposition rate than the concentrations in sediments.
- 25 Effects of invertebrate predators and a pesticide on temporary pond microcosms used for aquatic toxicity testing**
M.J. Barry, W. Davies
This study demonstrates that the use of temporary pond microcosms containing invertebrate predators may be a robust and useful method for ecosystem level toxicity testing.
- 35 Concentrations and gas/particle partitioning of PCBs in Chicago**
Y. Tasdemir, N. Vardar, M. Odabasi, T.M. Holsen
Results suggest that the area southwest of Chicago can be an important local or regional source sector for PCBs.
- 45 Treatment of atrazine by integrating photocatalytic and biological processes**
C.Y. Chan, S. Tao, R. Dawson, P.K. Wong
Photochemical pretreatment plus microbial degradation lead to complete degradation and detoxification of atrazine.
- 55 Variability of catechin and 4-hydroxyacetophenone distribution in Norway spruce needles in relation to their position, age, and growing conditions**
N. Vrhotová, J. Triska, O. Urban, L. Pěkníc
Catechin concentration in spruce needles was influenced by tree position and needle position in the tree canopy.
- 61 Solute transport and extraction by a single root in unsaturated soils: model development and experiment**
J. Kim, K. Sung, M.Y. Corapcioglu, M.C. Drew
The single-root model and experiment were useful in understanding microscopic phenomena of contaminant transport in soil-plant system.
- 71 Further investigation of the heavy metal content of the teeth of the bank vole as an exposure indicator of environmental pollution in Poland**
J. Gdula-Argasińska, J. Appleton, K. Sawicka-Kapusta, B. Spence
The molar teeth of bank voles are sensitive and accurate bio-indicators of heavy metals' exposure and reflect decreased metals deposition related to the improved environmental conditions.
- 81 A lysimeter experiment to investigate temporal changes in the availability of pesticide residues for leaching**
F.G. Renaud, C.D. Brown, C.J. Fryer, A. Walker
Leaching of aged herbicides varies with properties of the compound and of the soil.
- 93 Comparison of throughfall and soil solution chemistry between a high-density Corsican pine stand and a naturally regenerated silver birch stand**
A. De Schrijver, L. Nachtergale, J. Staelens, S. Luysaert, L. De Keersmaeker
Nitrogen throughfall deposition in a naturally regenerated birch stand was 3.5 times lower than in a neighbouring Corsican pine plantation.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; ISI GeoSciTech; Science Citation Index; SciSearch)




0269-7491(200409)131:1;1-Z



ELSEVIER

02022

Volume 131 No. 2 September 2004 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 131 Number 2 2004

- 773 Modelling the spatial distribution of ammonia emissions from seabirds in the UK**
L.J. Wilson, P.J. Bacon, J. Bull, U. Dragosits, T.D. Blackall, T.E. Dunn, K.C. Hamer, M.A. Sutton, S. Wanless
Ammonia emissions from each seabird species were estimated and combined with population distribution data.
- 187 Phytoavailability and fractionation of copper, manganese, and zinc in soil following application of two composts to four crops**
V.D. Zheljzkov, P.R. Warman
Results suggest that composted municipal solid waste with high Cu and Zn can be used safely with agricultural crops.
- 197 Trace metal and stable isotope measurements ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) in the harbour porpoise *Phocoena phocoena relicta* from the Black Sea**
K. Das, L. Holsbeek, J. Browning, U. Siebert, A. Birkun, Jr, J.-M. Bouqueneau
Trace metal levels in Black Sea harbour porpoises are lower than levels from the North Sea.
- 205 Does interspecific competition alter effects of early season ozone exposure on plants from wet grasslands? Results of a three-year experiment in open-top chambers**
A.E.G. Tonneijck, J. Franzaring, G. Brouwer, K. Metselaar, Th.A. Dueck
Interspecific competition between plants from wet grasslands does not alter effects of early season ozone on plant growth in a three-year experiment.
- 215 Effects of lead-contaminated sediment and nutrition on mallard duckling brain growth and biochemistry**
E. Douglas-Stroebel, D.J. Hoffman, G.L. Brewer, L. Sileo
Ingestion of lead-contaminated sediment and nutrition affects mallard duckling brain growth and biochemistry with respect to oxidative stress and other metabolic alterations.
- 223 Estimation of apparent rate coefficients for phenanthrene and pentachlorophenol interacting with sediments**
H.-L. Chen, Y.-X. Chen, Y.-T. Xu, M.-W. Shen
A model was used to study time-dependent interactions of phenanthrene and PCP in collected sediments.
- 233 Comparison of different physiological parameter responses in *Lemna minor* and *Scenedesmus obliquus* exposed to herbicide flumioxazin**
L. Geoffroy, C. Frankart, P. Eullaffroy
Interspecific variations and the nature of parameters both affected toxicity.
- 243 Metal availability and soil toxicity after repeated croppings of *Thlaspi caerulescens* in metal contaminated soils**
C. Keller, D. Hammer
*Repeated croppings with *Thlaspi caerulescens* reduce Cd and Zn availability and toxicity in an acidic soil but not in a calcareous one.*
- 255 Biosensors for detection of mercury in contaminated soils**
I. Bontidean, A. Mortari, S. Leth, N.L. Brown, U. Karlson, M.M. Larsen, J. Vangronsveld, P. Corbisier, E. Csöregi
Whole bacterial cell and protein-based biosensors were effective in assessing mercury in soil samples.
- 263 Biotransformation of 2,4,6-trinitrotoluene (TNT) by enchytraeids (*Enchytraeus albidus*) in vivo and in vitro**
S.G. Dodard, J. Powlowski, G.I. Sunahara
Enchytraeids transformed TNT in vivo and in vitro, with apparent involvement of bacteria associated with the host organism.
- 275 Mercury in fish from the Pinchi Lake Region, British Columbia, Canada**
S.A. Weech, A.M. Scheuhammer, J.E. Elliott, K.M. Cheng
Hg concentrations in small (<40 cm) fish from Pinchi Lake, British Columbia, Canada are elevated compared with those from nearby reference lakes.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; ISI GeoSciTech; Science Citation Index; SciSearch)




0269-7491(200409)131:2;1-Y



ELSEVIER

02022



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



CONTENTS

Volume 131 Number 3 2004

- 327 Thiol synthesis and arsenic hyperaccumulation in *Pteris vittata* (Chinese brake fern)**
W. Zhang, Y. Cai, K.R. Downum, L.Q. Ma
Results suggest that acid-soluble thiols may play a supplementary role in arsenic detoxification.
- 347 The effect of ozone on pollen development in *Lolium perenne* L.**
K. Schoene, J.-Th. Franz, G. Masuch
*Starch accumulation during pollen development of *Lolium perenne* was strongly inhibited after ozone exposure of plants.*
- 355 Evaluation of the BCR sequential extraction procedure applied for two unpolluted Spanish soils**
E. Fernández, R. Jiménez, A.M. Lallena, J. Aguilar
The sequential extraction procedure of BCR is suitable for As, Pb and Cd at low concentrations, but not for Cu, Co, Cr, Zn and Ni.
- 365 The effects of the contemporary-use insecticide (fipronil) in an estuarine mesocosm**
E.F. Wirth, P.L. Pennington, J.C. Lawton, M.E. DeLorenzo, D. Bearden, B. Shaddrix, S. Sivertsen, M.H. Fulton
Fipronil exposure at environmentally relevant concentrations had a significant toxic effect on grass shrimp in mesocosms.
- 373 Relationships between metal concentrations in great tit nestlings and their environment and food**
T. Dauwe, E. Janssens, L. Bervoets, R. Blust, M. Eens
Excreta of nestling passerines may be a suitable monitor for the presence and concentrations of metals in the food and the environment.
- 381 Persistent organochlorines in human breast milk collected from primiparae in Dalian and Shenyang, China**
T. Kunisue, M. Someya, F. Kayama, Y. Jin, S. Tanabe
The infants in Dalian may be exposed to relatively high levels of DDTs and HCHs through mother's milk.
- 393 Manganese uptake and accumulation by the hyperaccumulator plant *Phytolacca acinosa* Roxb. (Phytolaccaceae)**
S.G. Xue, Y.X. Chen, R.D. Reeves, A.J.M. Baker, Q. Lin, D.R. Fernando
*A newly discovered hyperaccumulator of manganese, *Phytolacca acinosa* Roxb., has phytoremediation potential.*
- 401 Concentrations of organotin compounds in sediment and clams collected from coastal areas in Vietnam**
S. Midorikawa, T. Arai, H. Harino, M. Ohji, N. Duc Cu, N. Miyazaki
Sediment-bound TBT is another source of contamination for clams in coastal areas of Vietnam.
- 409 Interactions of metals affect their distribution in tissues of *Phragmites australis***
J.S. Weis, T. Glover, P. Weis
**Phragmites australis*, when grown with two or more metals, alters its distribution of a particular metal from the distribution after exposure to a single metal, resulting usually in greater aboveground allocation.*
- 417 Changes in the atmospheric deposition of minor and rare elements between 1975 and 2000 in south Sweden, as measured by moss analysis**
Å. Rühling, G. Tyler
Reductions in anthropogenic dust emissions has reduced deposition of most elements in the periodic table.
- 425 Organochlorine pesticides and polychlorinated biphenyls in California sea lions**
K. Kannan, N. Kajiwara, B.J. Le Boeuf, S. Tanabe
PCB and DDT concentrations in California sea lions continue to be of concern.
- 435 Mechanisms of lead, copper, and zinc retention by phosphate rock**
X. Cao, L.Q. Ma, D.R. Rhue, C.S. Appel
Phosphate-induced formation of fluoropyromorphite is primarily responsible for Pb immobilization by phosphate rock, whereas Cu and Zn retention is mainly attributable to the surface adsorption or complexation.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; ISI GeoSciTech; Science Citation Index; SciSearch)



0269-7491(200410)131:3;1-2



ELSEVIER

02022